

Topology
of
non-topological Chern-Simons vortices

P. A. HORVÁTHY

Laboratoire de Mathématiques et de Physique Théorique

Université de Tours

Parc de Grandmont, F-37200 TOURS (France) ⁽¹⁾

Abstract. *The quantized magnetic flux $\Phi = -4\pi N(\text{sign } \kappa)$, $N = 0, \pm 1, \dots$ of non-topological vortices in the non-relativistic Chern-Simons theory is related to the topological degree of the $S^2 \rightarrow S^2$ mapping defined by lifting the problem to the Riemann spheres. Regular solutions with finite degree only arise for rational functions, whose topological degree, N , is the common number of their zeros and poles on the Riemann sphere, also called their algebraic order.*

(1/2/2008)

PACS numbers: 0365.GE, 11.10.Lm, 11.15.-q

⁽¹⁾ e-mail: horvathy@univ-tours.fr

A striking feature of non-relativistic Chern-Simons vortices [1] is the quantization of their magnetic flux,

$$(1) \quad \text{vortex flux} = \int_{\mathbf{R}^2} B d^2\vec{x} = -4\pi N(\text{sign } \kappa), \quad N = 0, \pm 1, \dots$$

In gauge theories, similar properties are usually related to *topology*. Yang–Mills–Higgs theories ⁽¹⁾, for example, support finite–energy monopole solutions [2]. For large r , the Higgs field is required to satisfy $\text{Tr}(\Phi)^2 = 1$ rather than to vanish. This asymptotic condition allows us to view Φ as a mapping from the two-sphere at infinity, $\mathbf{S} \equiv \mathbf{S}_\infty^2$, into the vacuum manifold $\text{Tr}(\Phi)^2 = 1$, which is again a two-sphere. Then the magnetic charge of the monopole is expressed [3] as

$$(2) \quad \text{Monopole charge} = \frac{1}{8\pi} \int_{\mathbf{S}} \epsilon_{ijk} \epsilon_{abc} \Phi^a \partial_j \Phi^b \partial_k \Phi^c d^2 S.$$

The r. h. s. here is a topological invariant characterized by an integer. It is in fact the homotopy class (or “winding number”) of Φ , $[\Phi] \in \pi_2(\mathbf{S}^2) \simeq \mathbf{Z}$.

Non-relativistic Chern-Simons vortices are, however, *non-topological*: for finite–energy solutions the particle density, ϱ , tends to zero when spatial infinity is approached. Then where does flux quantization, Eq. (1), come from ?

To answer this question, let us remember that, owing to the “field-current identity” which relates the magnetic field to the particle density, $\kappa B = -\varrho$, the magnetic flux of the vortex is

$$(3) \quad \text{vortex flux} = -\frac{1}{\kappa} \int_{\mathbf{R}^2} \varrho d^2\vec{x}.$$

Here, apart of the zeros, ϱ is a solution of the Liouville equation $\Delta \log \varrho = -(2/|\kappa|)\varrho$. Now the general solution of the Liouville equation is expressed using an arbitrary analytic function $f(z)$,

$$(4) \quad \varrho = 4|\kappa| \frac{|f'|^2}{(1 + |f|^2)^2},$$

so that

$$(5) \quad -\frac{(\text{sign } \kappa)}{4\pi} (\text{vortex flux}) = \frac{1}{4\pi} \int_{\mathbf{R}^2} \frac{4|f'|^2}{(1 + |f|^2)^2} dx dy.$$

The point is that the r. h. s. has the *same* topological interpretation as (2) for monopoles. The analytic function f can in fact be viewed as mapping the z -plane into the w -plane. Compactifying these planes into Riemann spheres $\mathbf{S}_z \simeq \mathbf{C} \cup \{\infty\}$ and $\mathbf{S}_w \simeq \mathbf{C} \cup \{\infty\}$,

⁽¹⁾ We assume, for simplicity, that the gauge group is SU(2).

z and w become stereographic coordinates. If f admits a well-defined (finite or infinite) limit as $r \rightarrow \infty$, it can be extended into a mapping from $\mathbf{S} \equiv \mathbf{S}_z$ into $\mathbf{S}_w$ we still denote by f . The extended mapping has a winding number, given in (2). This latter is in fact the integral of the pull-back by f of Ω , the surface form of the unit sphere $\mathbf{S}_w$,

$$(6) \quad \text{charge} = \frac{1}{4\pi} \int_{\mathbf{S}} f^* \Omega.$$

Now, in complex coordinate w , the surface element of $\mathbf{S}_w$ is

$$(7) \quad \Omega = 2i \frac{dw \wedge d\bar{w}}{(1 + w\bar{w})^2},$$

so that setting $w = f(z)$ in Eq. (6) yields precisely the r. h. s. of (5), as stated.

In a previous paper [4] we proved that the regularity of finite-flux solutions excludes essential singularities in the finite domain as well as at infinity. An analytic function whose only singularities in the extended plane are poles is known to be a *rational function*,

$$(8) \quad f(z) = \frac{a_m z^m + \dots + a_1 z + a_0}{b_n z^n + \dots + b_1 z + b_0}.$$

[5]. But a rational function has always a limit at infinity, namely $\infty \in \mathbf{S}_w$ if $m > n$, $0 \in \mathbf{S}_w$ if $m < n$ and $a_m/b_m \neq 0$, ∞ if $m = n$. In the first case, $\infty \in \mathbf{S}_z$ is a pole of order $m - n$ while in the second it is a zero of order $n - m$. Since the only other poles and zeros are the n zeros of the denominator and the m zeros of the numerator, respectively, we see that the extended map $f : \mathbf{S}_z \rightarrow \mathbf{S}_w$ has the same number of zeros and poles, namely $N = \max(m, n)$. This common number is called the *algebraic order* of the rational map f [5].

The magnetic charge (2) has another useful description [3]. The r. h. s. of (2) is in fact equal to the *topological* (or Brouwer) *degree* of the field Φ . Choosing coordinates on the spheres, the mapping Φ can be characterized by $\Phi^a(\xi)$, $\xi = (\xi^b)$ $a, b = 1, 2$. A point Φ_0 in the image-sphere is a regular point if $\det(\partial\Phi^a/\partial\xi^b) \neq 0$. Then the topological degree is the number of pre-images of a regular point Φ_0

$$(9) \quad \text{degree} = \sum_{\xi \in \Phi_0} \text{sign} \det \left(\frac{\partial \Phi^a}{\partial \xi^b} \right).$$

Eq. (9) counts the number of times the image-two-sphere is covered while the domain-two-sphere is covered once. Then, assuming that ∞ and 0 are regular points, the topological degree of the rational function (8) is simply the number of zeros or, equivalently, the number of poles. Thus, we have proved

Theorem1. *The magnetic flux of the vortex solution associated with the rational function f is (1), where N is the algebraic order of f .*

The same result was obtained in Ref. [4] by a rather laborious direct calculation.

Let us also remark that the vortex number could also be identified with the first Chern class of an $U(1)$ principal bundle over the two-sphere.

The correspondence between analytic functions f and solutions ϱ of the Liouville equation is not one-to-one, though. Let R denote an arbitrary rotation of the image-two-sphere $\mathbf{S}_w$. Then f and $R \circ f$ yield the same solutions. This is seen by noting that the particle density is in fact the dual on the two-sphere $\mathbf{S} \equiv \mathbf{S}_z$ of the two-form $f^* \Omega$, $\varrho = \star(f^* \Omega)$. But

$$(R \circ f)^* \Omega = f^*(R^* \Omega) = f^* \Omega,$$

because the surface-form Ω is symmetric w. r. t. rotations, $R^* \Omega = \Omega$. There is hence no loss of generality in requiring that

$$(10) \quad f(\infty) = 0,$$

since this can always be achieved by a rotation of $\mathbf{S}_w$. A rational function which satisfies the condition (10) can be written as

$$(11) \quad f(z) = \frac{P(z)}{Q(z)}, \quad \deg P < \deg Q$$

whose order is $N = \deg Q$. Such an f depends on the $2N$ complex parameters, namely the coefficients of the polynomials $P(z)$ and $Q(z)$. Thus, we have proved

Theorem2. *For fixed magnetic flux, the solution depends on $4N$ real parameters.*

Requiring $f(\infty) = \infty$ instead of (10), the role of poles and zeros would be interchanged, without changing the conclusion.

Theorem2 is again consistent with previous results, [4], [6]. The particular form (11) was reached in Ref. [4] by exploiting the invariance of the solution by $f \rightarrow (1/f)$ when $m > n$ in (8), and by reducing the case $m = n$ to (11) by a suitable redefinition. These transformations correspond simply to $O(3)$ transformations which interchange poles and zeros or rotate a point $\neq 0, \infty$ into the origin, respectively.

Let us note, in conclusion, that our result also applies to the $O(3)$ sigma model which describes a two-dimensional ferromagnet [7]. The scalar field Φ is here a mapping from the plane into a two-sphere $\mathbf{S}^2$. For large distances, all magnets are required to be aligned so that $\Phi(\infty) = \Phi_0 \in \mathbf{S}^2$. This allows to extend Φ into a mapping between two-spheres. The self-dual solutions found by Belavin and Polyakov [7] are associated with analytic functions on the complex plane with identical properties as here above.

Acknowledgement. It is a pleasure to thank N. Manton and C. Duval for illuminating discussions.

References

- [1] R. Jackiw and S-Y. Pi, *Phys. Rev. Lett.* **64**, 2969 (1990); *Phys. Rev.* **D42**, 3500 (1990). For review see R. Jackiw and S-Y. Pi, *Prog. Theor. Phys. Suppl.* **107**, 1 (1992), or G. Dunne, *Self-Dual Chern-Simons Theories*. Springer Lecture Notes in Physics. New Series: Monograph 36. (1995).
- [2] For a review on monopoles see, for instance, P. Goddard and D. Olive, *Monopoles in gauge theory*, *Rep. Prog. Phys.* **41**, 1357-1437 (1978).
- [3] J. Arafune, P G. O. Freund, and C. J. Goebel, *Journ. Math. Phys.* **16**, 433 (1975). The topological facts used in this paper are taken from J. W. Milnor, *Topology from the Differentiable Viewpoint*, Univ. of Virginia Press, Charlottesville (1965).
- [4] P. A. Horváthy and J.-C. Yéra, *Lett. Math. Phys.* **46**, 111-120 (1998).
- [5] L. V. Ahlfors, *Complex analysis*, McGraw–Hill (1979). See also E. T. Whittaker and G. N. Watson, *A course of modern analysis*. Fourth edition. Cambridge University Press. Reprinted in 1992.
- [6] S. K. Kim, K. S. Soh, and J. H. Yee, *Phys. Rev.* **D42**, 4139 (1990).
- [7] A. A. Belavin, A. M. Polyakov, A. S. Schwartz and Yu. S. Tyupkin, *Phys. Lett.* **59B**, 85 (1975). The hypersphere description was proposed by R. Jackiw and C. Rebbi, *Phys. Rev.* **D14**, 517 (1976).
- [8] A. A. Belavin and A. M. Polyakov, *JETP. Lett.* **22**, 245 (1975).